


Arthritis of the Base of the Thumb

What is it?

In a normal joint, cartilage covers the end of the bones and serves as a shock absorber to allow smooth, pain-free movement. In osteoarthritis (OA, or “degenerative arthritis”) the cartilage layer wears out, resulting in direct contact between the bones and producing pain and deformity. One of the most common joints to develop OA in the hand is the base of the thumb. The thumb basal joint, also called the carpometacarpal (CMC) joint, is a specialized saddle-shaped joint that is formed by a small bone of the wrist (trapezium) and the first bone of the thumb (metacarpal). The saddle shaped joint allows the thumb to have a wide range of motions, including up, down, across the palm, and the ability to pinch (see Figure 1).




Who gets it?

OA at the base of the thumb is more commonly seen in women over the age of 40. The exact cause is unknown, but genetics, previous injuries such as fractures or dislocations, and generalized joint laxity may predispose towards development of this type of arthritis.

What are the symptoms and signs?

The most common symptom is pain at the base of the thumb. The pain can be aggravated by activities that require pinching, such as opening jars, turning door knobs or keys, and writing. Severity can also progress to pain at rest and pain at night. In more severe cases, progressive destruction and mal-alignment of the joint occurs, and a bump develops at the base of the thumb as the metacarpal moves out of the saddle joint. This shift in the joint can cause limited motion and weakness, making pinch difficult (see Figure 2). The next joint above the CMC may compensate by loosening, causing it to bend further back (hyperextension).




How is the diagnosis made?

The diagnosis is made by history and physical evaluation. Pressure and movement such as twisting will produce pain at the joint. A grinding sensation may also be present at the joint (see Figure 3). X-rays are used to confirm the diagnosis, although symptom severity often does not correlate with x-ray findings.

What are the treatment options?

Less severe thumb arthritis will usually respond to non-surgical care. Arthritis medication, splinting and limited cortisone injections may help alleviate pain. A hand therapist might provide a variety of rigid and non-rigid splints which can be used while sleeping or during activities.

Patients with advanced disease or who fail non-surgical treatment may be candidates for surgical reconstruction. A variety of surgical techniques are available that can successfully reduce or eliminate pain. Surgical procedures include removal of arthritic bone and joint reconstruction (arthroplasty), joint fusion, bone realignment, and even arthroscopy in select cases. A consultation with your hand surgeon can help decide the best option for you (see Figure 4).





Arthritis of the Thumb

Arthritis is a condition that irritates or destroys a joint. Although there are several types of arthritis, the one that most often affects the joint at the base of the thumb (the basal joint) is osteoarthritis (degenerative or "wear-and-tear" arthritis).

Description

Smooth cartilage covers the ends of the bones. It enables the bones to glide easily in the joint. Without it, bones rub against each other, causing friction and damage to the bones and the joint. Osteoarthritis occurs when the cartilage begins to wear away.

The joint at the base of the thumb, near the wrist and at the fleshy part of the thumb, enables the thumb to swivel, pivot, and pinch so that you can grip things in your hand.

Arthritis of the base of the thumb is more common in women than in men, and usually occurs after 40 years of age. Prior fractures or other injuries to the joint may increase the likelihood of developing this condition.



Symptoms

- Pain with activities that involve gripping or pinching, such as turning a key, opening a door, or snapping your fingers
- Swelling and tenderness at the base of the thumb
- An aching discomfort after prolonged use
- Loss of strength in gripping or pinching activities
- An enlarged, "out-of-joint" appearance
- Development of a bony prominence or bump over the joint
- Limited motion

Doctor Examination

Your doctor will ask you about your symptoms, any prior injury, pain patterns, or activities that aggravate the condition.

One of the tests used during the examination involves holding the joint firmly while moving the thumb. If pain or a gritty feeling results, or if a grinding sound (crepitus) can be heard, the bones are rubbing directly against each other.

An x-ray may show deterioration of the joint as well as any bone spurs or calcium deposits that have developed.

Many people with arthritis at the base of the thumb also have symptoms of carpal tunnel syndrome, so your doctor may check for that, as well.

Treatment

Nonsurgical Treatment

In its early stages, arthritis at the base of the thumb will respond to nonsurgical treatment.

- Ice the joint for 5 to 15 minutes several times a day.
- Take an anti-inflammatory medication, such as aspirin or ibuprofen, to help reduce inflammation and swelling.
- Wear a supportive splint to limit the movement of your thumb, and allow the joint to rest and heal. The splint may protect both the wrist and the thumb. It may be worn overnight or intermittently during the day.

Because arthritis is a progressive, degenerative disease, the condition may worsen over time. The next phase in treatment involves injecting a steroid solution injection directly into the joint. This will usually provide relief for several months. However, these injections cannot be repeated indefinitely.

Surgical Treatment

When nonsurgical treatment is no longer effective, surgery is an option. The operation can be performed on an outpatient basis, and several different procedures can be used.

One option involves fusing the bones of the joint together. This, however, will limit movement.

Another option is to remove part of the joint and reconstruct it using either a tendon graft or an artificial substance.

You and your physician will discuss the options and select the one that is best for you.

Rehabilitation

After surgery, you will have to wear a cast for 4 to 8 weeks, depending on which procedure is used. A rehabilitation program, often involving a physical therapist, helps you regain movement and strength in the hand. You may feel some discomfort during the initial stages of the rehabilitation program, but this will diminish over time. Full recovery from surgery takes several months. Most patients are able to resume normal activities.

Last reviewed: December 2013

AAOS does not endorse any treatments, procedures, products, or physicians referenced herein. This information is provided as an educational service and is not intended to serve as medical advice. Anyone seeking specific orthopaedic advice or assistance should consult his or her orthopaedic surgeon, or locate one in your area through the AAOS ["Find an Orthopaedist"](#) program on this website.

Copyright 2013 American Academy of Orthopaedic Surgeons

Related Topics

Arthritis: An Overview (<http://orthoinfo.aaos.org/topic.cfm?topic=A00208>)

Arthritis of the Hand (<http://orthoinfo.aaos.org/topic.cfm?topic=A00224>)

What are NSAIDs? (<http://orthoinfo.aaos.org/topic.cfm?topic=A00284>)

OrthoInfo

The American Academy of Orthopaedic Surgeons
9400 West Higgins Road
Rosemont, IL 60018
Phone: 847.823.7186
Email: orthoinfo@aaos.org



SURGERY

Thumb Base Surgery

What does this involve?

Surgery on the base of the thumb involved removing some, or all, of the arthritic joint and replacing it with something less painful. Many different operations have been described for this but none has been conclusively shown to be better than any of the others. The simplest operations have the fewest complications, however.

When is surgery needed?

This operation is performed to relieve pain in an arthritic joint. Most surgeons would encourage you to try painkillers, splints, activity modification, aids to help with certain tasks (opening jars etc) and steroid injections into the painful joint before recommending surgery. The majority of patients with thumb base arthritis can manage their symptoms in this way without ever needing an operation.

Which operation is the right one for me?

The standard operation for this condition is a simple trapeziumectomy. In this operation the trapezium bone is removed preserving the capsule around the joint is tightened up to keep the thumb well aligned and a moulded plaster is applied for 4 weeks. Sometimes a wire is also used to support the bones whilst the soft tissues heal – this is removed later in clinic. During the time in plaster the space where the trapezium was fills in with scar tissue which acts as a soft pad for the thumb to work against. You then need to work on regaining movement of the thumb, which usually takes another 4 to 6 weeks.

If the tissue around the base of the thumb is very lax a trapeziumectomy may be combined with a ligament reconstruction and tendon interposition. The ligament reconstruction is usually done with part of a tendon from the front of your wrist passed through the base of the thumb metacarpal and sewn back on itself to act as a

sling around the thumb base. The end of the tendon is then folded up and put into the space where the trapezium was to act as extra padding.

Various other sorts of 'replacement' have been tried for this joint. Solid joints put into the small bones (rather like a tiny hip replacement) have been very difficult to fix in place and often loosen after a short period of time and cause more problems. More recently interest has turned to using a solid, smooth ring of pyrocarbon (a material which has been used for many years in heart valve surgery) as a replacement (The Pyrodisk). The end of the trapezium and base of the thumb metacarpal are removed and the circular replacement put between the bone surfaces. A ligament reconstruction is then carried out but the strip of tendon passes through the remains of the trapezium, the replacement and the base of the thumb metacarpal.

If very heavy loading of the thumb is likely to be necessary (eg working in the building profession etc) a fusion of the thumb base joint may be the preferred option. Your surgeon will discuss the best option with you.

X-Ray of left hand showing thumb base arthritis



Type of Operation

Day case

Length of Procedure

1 hour

Anaesthesia

Regional Anaesthetic (whole arm numb) or General Anaesthetic (asleep)



What are the main risks of this operation?

Swelling, Stiffness and Scar pain

This can be reduced by keeping the arm elevated and moving all the free joints as soon as possible. In most people the general swelling reduces dramatically in the first week after the operation.

Local swelling around the surgical site can persist for several months. Local swelling can be helped by massaging the tissues and this may also improve any irritability in the surgical scar.

Occasionally patients are troubled by more swelling and stiffness than average. In this case Complex Regional Pain Syndrome (CRPS) is sometimes the cause (see relevant information sheet in 'Conditions we Treat'). Severe CRPS occurs in less than 1% of cases.

Infection

This is unusual in the hand (less than 1% of cases). Local wound infections can often be treated with oral antibiotics. Rare, deep seated infections may require re-admission to hospital, antibiotics into the vein and occasionally more surgery.

Nerve Damage

The nerves most at risk with these operations are the branches of the superficial radial nerve (supplying sensation on the back of the thumb, index and middle fingers) and the palmar branch of the median nerve (supplying skin sensation at the base of the front of the thumb). This second nerve is only at risk with ligament reconstruction operations. The lost patch of skin sensation from these injuries might be irritating but should not affect how your hand works. Damage to these nerves can sometimes make CRPS worse.

Worsening of Carpal Tunnel Syndrome

Patients with evidence of carpal tunnel syndrome before thumb base surgery may find that these symptoms are much worse around the time of the operation, probably because of increased swelling in the area. Many surgeons would, therefore, consider decompressing the carpal tunnel at the same time as the thumb base surgery if you already have some signs of carpal tunnel syndrome.

Residual thumb base discomfort

No operation for thumb base arthritis is perfect. Some patients will still have some discomfort at the base of the thumb with any of the operations described above. In most cases the pre-operative pain is much reduced and the residual symptoms are a manageable nuisance only. Occasionally the symptoms are so marked as to require further surgery. For instance the base of the thumb may become unstable after a simple trapeziumectomy necessitating a ligament reconstruction or the joints left behind after a Pyrodisk has been inserted may wear out and need more attention.

Reduced pinch grip strength (thumb to index finger)

This is likely to be less than in a normal thumb after these operations but more than before the operation, when the joint was very painful. One of the goals of solid replacements or a fusion is to gain more pinch grip strength than with a soft tissue procedure.

Post Operative Course

Day 1-14

- A dressing and padded bandage with a splint or plaster support incorporated is applied after the operation
- Keep the dressings clean and dry
- Keep the arm elevated in a sling or on pillows to reduce swelling
- Start moving all the joints that are not immobilised as soon as possible to prevent stiffness (see Finger exercise sheet in 'Rehabilitation')
- Take painkillers before the anaesthetic wears off and as necessary thereafter

Two Weeks

- An appointment will be made for a wound check, removal of stitches (if necessary) and a new moulded cast to be applied

4 Weeks

- Plaster will be removed and rehabilitation will start from a removable splint under the supervision of the hand therapists

6 Weeks

- Strengthening starts
- Splint only being worn at night

3 Months

- Contact sports may be re-started

Plaster Cast Information

Contact your surgical centre if:

- Your fingers become blue, swollen or numb and tingling with a plaster cast in place
- You see any discharge, wetness or detect any unpleasant smells from around your cast
- The cast becomes cracked, soft, loose or uncomfortable.

Outside normal working hours you may need to attend your local Accident and Emergency Department for help with these issues.

Driving

You may drive when you feel confident to control the car, even in an emergency.

For thumb base surgery it is usually two months before you should consider driving again.

You should discuss it with your insurer if you are considering driving with a splint in place.

Time off Work

This will vary depending on the nature of your job and the exact nature of your surgery.

Sick notes can be provided on the day of your operation, at your clinic visits and by your own GP.

These notes are intended as a guide and some of the details may vary depending on your individual circumstance and at the discretion of your surgeon.